


DATA SHEET

Product Information

Product Name	: Recombinant Human Interleukin 2 (rhIL-2)
Synonyms	: IL2, TCGF, lymphokine, interleukin 2
Description	: Interleukin-2 (IL-2) is an interleukin, a type of cytokine signaling molecule in the immune system. It is a protein that regulates the activities of white blood cells that are responsible for immunity. In the thymus, where T cells mature, it prevents autoimmune diseases by promoting the differentiation of certain immature T cells into regulatory T cells, which suppress other T cells that are otherwise primed to attack normal healthy cells in the body. IL-2 also promotes the differentiation of T cells into effector T cells and into memory T cells when the initial T cell is also stimulated by an antigen, thus helping the body fight off infections. (Ref. From Wikipedia)
NCBI Accession No.	: NM_000586.3
Amino acid sequence	: MAPTSSSTKKTQLQLEHLLLDLQMI L N G I N N Y K N P K L T R M L T F K F Y M P K K A T E L K H L Q C L E E E L K P L E E V L N L A Q S K N F H L R P R D L I S N I N V I V L E L K G S E T T F M C E Y A D E T A T I V E F L N R W I T F C Q S I I S T L T
Molecular Mass	: 15.5 kDa (134 a.a)
Protein Tags	: No tagging
Source	: <i>E. coli</i> .
Cat. No.	: JW-H031-0010, JW- H031-0025, JW-H031-0100, JW-H031-0250, JW-H031-0500, JW-H031-1000
Storage	: Should be at ≤ -70 °C as undiluted aliquots of handy size. Avoid repeated freezing and thawing.
Cross Reactivity	: Human, Mouse, Rat, Monkey

Quality Control

Test items	Specifications
Appearance	: Clear, colorless liquid
Purity	: Greater than 98 % by RP-HPLC and SDS-PAGE
Specificity	: Using Western blot, detection
Concentration	: 0.1 mg/ml, Bradford method Determined by proliferation of CTLL-2 indicator cells.
Biological Activity	: The ED50 is ≤ 0.25 ng/ml, corresponding to a specific activity of $\geq 4.0 \times 10^6$ IU/mg
Endotoxin	: Less than 1.0 EU/ μ g as determined by the LAL method
Formulation	: 50mM Tris, 10% Glycerol (pH 8.0) without preservative or carrier proteins.
Stability	: Stable for up to 12 months at -70 °C. Stable for a month at 4 °C.
Sterility	: Sterilized through a 0.2 μ m membrane filter and packaged aseptically. Culture for 2 weeks, no growth



Manufacturer : **JW CreaGene Inc.**
 FL2, Jungang induspia V, 138-6, Sangdaewon-dong, Jungwon-gu, Seongnam-si,
 Gyeonggi-do, South Korea (462-120)
 Tel: 82-31-737-3310, Fax: 82-31-737-3301