

DATA SHEET

Product Name	:	Recombinant Mouse Interleukin-1 Beta (rmIL-1β)
Synonyms	:	IL-1 beta, Catabolin, <i>IL1B</i>
Description	:	Interleukin-1 beta (IL-1 β) known as catabolin, is a cytokine protein that in humans is encoded by the IL1B gene. IL-1 β precursor is cleaved by cytosolic caspase 1 (interleukin 1 beta convertase) to form mature IL- 1 β . Interleukin 1 was discovered by Igal Gery in 1972. He named it lymphocyte-activating factor (LAF) because it was a lymphocyte mitogen. It was not until 1985 that interleukin 1 was discovered to consist of two distinct proteins, now called interleukin-1 alpha and interleukin-1 beta. IL-1 β is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. (Ref. From Wikipedia)
NCBI Accession No.	:	NM_008361.4
Amino acid sequence	:	MVPIRQLHYRLRDEQQKSLVLSDPYELKALHLNGQNINQQVIFSMSFVQGEP SNDKIPVALGLKGKNLYLSCVMKDGTPTLQLESVDPKQYPKKKMEKRFVFNK IEVKSKVEFESAEFPNWYISTSQAEHKPVFLGNNSGQDIIDFTMESVSS*
Molecular Mass	:	17.52 kDa (153 aa)
Protein Tags	:	No tagging
Source	:	E. coli.
Cat. No.	:	JW-M007-0010, JW-M007-0025, JW-M007-0100, JW-M007-0250, JW- M007-0500, JW-M007-1000
Storage	:	Should be at \leq -70 °C as undiluted aliquots of handy size. Avoid repeated freezing and thewing
Cross Reactivity	:	repeated freezing and thawing. Human, Monkey, Rat, Pig, Chicken

Quality Control

Test items	Specifications
Appearance	: Clear, colorless liquid
Purity	: Greater than 98 % by RP-HPLC and SDS-PAGE
Specificity	: Using Western blot, detection TF-1 proliferation assay [2X10 ⁴ : rmlL-1beta, 48h]
Concentration	: 0.1 mg/ml, Bradford method
Biological Activity	Determined by proliferation of TF-1 cells : The ED50 is < 10 pg/ml, corresponding to a specific activity of > 1.0 × 10 ⁸ U/mg
Endotoxin	: Less than 0.1 EU/ μ g as determined by the LAL method
Formulation	: PBS, 5% glycerol (pH 7.4) without preservative or carrier proteins.
Stability	: Stable for up to 12 months at -70 °C. Stable for a month at 4 °C.
Sterility	Sterilized through a 0.2 μm membrane filter and packaged aseptically. Culture for 2 weeks, no growth

Manufacturer

FL2, Jungang induspia V, 138-6, Sangdaewon-dong, Jungwon-gu, Seongnam-si, Gyeonggi-do, South Korea (462-120) Tel: 82-31-737-3310, Fax: 82-31-737-3301